

UNCLASSIFIED

AD NUMBER
ADB247697
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info; Oct 98, U.S. Army Medical Research and Materiel Command, 504 Scott Street, Other requests shall be referred to , Ft Detrick, MD 21702-5012
AUTHORITY
USAMRMC ltr, 23 Aug 2001

THIS PAGE IS UNCLASSIFIED

AD_____

GRANT NUMBER DAMD17-97-1-7302

TITLE: A Comparison of Breast Cancer Treatment Regimens by
Demographic Characteristics

PRINCIPAL INVESTIGATOR: Marianne E. Ulcickas Yood, M.P.H.

CONTRACTING ORGANIZATION: Henry Ford Health System
Detroit, Michigan 48202-3450

REPORT DATE: October 1998

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S.
Government agencies only (proprietary information, October
1998). Other requests for this document shall be referred
to U.S. Army Medical Research and Materiel Command, 504
Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

19990928 409

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1998		3. REPORT TYPE AND DATES COVERED Annual (5 Sep 97 - 4 Sep 98)
4. TITLE AND SUBTITLE A Comparison of Breast Cancer Treatment Regimens by Demographic Characteristics			5. FUNDING NUMBERS DAMD17-97-1-7302	
6. AUTHOR(S) Marianne E. Ulcickas Yood, M.P.H.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Henry Ford Health System Detroit, Michigan 48202-3450			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, October 1998). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>The purpose of this study was to measure ethnic differences in treatment and survival between African American (AA) and European American (EA) women with breast cancer. This annual report presents results of the survival analysis.</p> <p>We abstracted data on cases of breast cancer diagnosed in members of an HMO in metropolitan Detroit between 1986-1996 (N=886) and followed these cases for survival through April 1997 (N=137 deaths).</p> <p>AA women were diagnosed at a later stage when compared with EA women. Five-year survival was 77% for AAs and 84% for EAs. The crude hazard for AAs relative to EAs was 1.6 (95% confidence interval (CI) 1.1, 2.2). Adjusting only for stage of disease at diagnosis, the hazard ratio was 1.3 (95% CI 0.9, 1.9). Adjusting only for sociodemographics (age, marital status and income), the hazard ratio was 1.2 (95% CI 0.8, 1.9). After adjusting for sociodemographics and stage, the hazard ratio was 1.0 (95% CI 0.7, 1.5).</p> <p>Among women with similar medical care access, we found ethnic differences in stage of breast cancer at diagnosis. Adjusting for this difference and for income, age and marital status, eliminates the effect of race on survival.</p>				
14. SUBJECT TERMS Breast Cancer			15. NUMBER OF PAGES 19	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified
				20. LIMITATION OF ABSTRACT Limited

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


____ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

my ✓ ____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.



PI - Signature

10/2/98

Date

**A Comparison of Breast Cancer Treatment Regimens by Demographic
Characteristics (DAMD17-97-1-7302), Annual Report
October, 1998**

TABLE OF CONTENTS

Introduction.....	5
Body.....	6
Methods.....	6
Results.....	9
Conclusions.....	10
References.....	12
Table 1. Baseline Demographics and Tumor Characteristics.....	16
Table 2. Effect of Demographic and Tumor Characteristics on Survival Estimates....	17
Figure 1. Crude Kaplan-Meier Survival Estimates, by Race.....	18
Figure 2. Survival by Race, Adjusted for Age, Income & Marital Status and Stage....	19

INTRODUCTION

In the United States, survival for African American women with breast cancer is inferior to that for European American women. The 1970s and 1980s marked a time of relatively stable rates of mortality among European American women with breast cancer, but increasing rates for African Americans¹. However, the decline in mortality observed in the early 1990s for European American with breast cancer was not observed in African Americans^{1,2}. Poorer survival among African Americans has been attributed to biological characteristics of the tumor, advanced stage at diagnosis, lower socioeconomic status (SES), barriers to health care, diagnostic and treatment delays^{3,4} and a higher prevalence of comorbid conditions^{5,6}. Although use of mammography by African American women has been reported to lag behind Caucasian women⁷, recent research indicates that the racial discrepancy is narrowing⁸. However, it is too soon to see how increased use of mammography among African Americans will affect survival.

While most investigations have found variability in tumor histology at disease presentation across ethnic groups⁹⁻¹¹ and a few have not¹², researchers suggest that the disparity is related more to SES and its impact on diagnostic delays or even a lag in benefitting from medical advancements¹³, as opposed to inherent biologic differences. In most studies, use of multivariable models to control for differences in tumor biology and sociodemographic characteristics have reduced but not eliminated the racial differential in survival^{6,14-17}. Some studies have attributed the mortality differences to racial disparity in socioeconomic status, with biology playing a lesser role¹⁸⁻²¹.

We present analyses of breast cancer survival in a population of HMO members where the screening, diagnosis and treatment patterns are based on practice standards and are similar for all members of the population served within a large, multidisciplinary group practice. We selected this population to minimize heterogeneity in care delivery and to eliminate issues of financial barriers to health care.

BODY

Methods

Setting

The setting for this study was the Health Alliance Plan (HAP) health maintenance organization. HAP is located in Southeastern Michigan and is the largest health maintenance organization in Michigan, with more than 450,000 members. Approximately 20% of these members are African American, 53% are female, and 57% are cared for by physicians in the Henry Ford Medical Group (HFMG). Our study population was drawn from HAP members served by the HFMG. The HFMG is a large multispecialty group practice consisting of a hospital-based clinic in a large urban teaching hospital in Detroit (Henry Ford Hospital) and 26 satellites throughout Southeastern Michigan.

HFHS maintains a computerized tumor registry data base accredited by the American College of Surgeons. Registry staff use a thorough case finding system, including review of all pathology and cytology reports, as well as radiation and oncology consultations. The American Joint Commission on Cancer (AJCC) system is used to determine stage of disease by evaluating tumor size, extent of invasion, involvement of lymph nodes and presence of metastasises.

HFMG Registry staff link these data with the Detroit area Surveillance, Epidemiology and End Results (SEER) program, and conducts annual follow-up for vital status and recurrence. The annual follow-up rate is estimated at 94%.

Ascertainment of Cases

Using the HFMG cancer registry, we identified all African American and European American women with newly diagnosed incident breast cancer initially treated at HFMG from January 1986 through April 1996. To minimize heterogeneity in clinical practice and access to care before diagnosis, we limited the study population to women continuously enrolled in HAP for at least one year before diagnosis and assigned to a primary care physician within the HFMG at the time of diagnosis. We defined continuous enrollment as no more than a 60-day gap in coverage according to membership files.

Outcome Data

We used a several sources to identify follow-up data. First, we obtained vital status, date of death (if applicable) and date last known alive from the HFMG tumor registry. Next, for those women known to be alive, we used HFMG administrative billing data to obtain information about hospitalizations and outpatient visits from January 1986 through April 1997. We used the billing data to update the tumor registry date where appropriate.

Identification of Confounding Variables

Using the tumor registry, we obtained information on tumor characteristics (stage and tumor size) and demographics (date of birth, sex). We geocoded addresses from billing files into

census block groups. We imputed household income to each woman using median household income for block group from the 1990 census data. Information about duration of HAP membership after diagnosis and mammography benefits was downloaded from the HMO membership files.

Statistical Methods.

Before analysis, we categorized age at diagnosis (<55 years vs. \geq 55 years), marital status (unmarried, married) and income (imputed household income <\$35,000, \geq \$35,000). Stage of disease was selected as the variable best describing disease status. We also examined length of enrollment in HAP before diagnosis (<5 years, \geq 5 years). These variables were chosen based on known relationships with both breast cancer survival and race (i.e., as potential confounders) rather than through statistical methods for selection of variables.

Association with survival was assessed using the hazard ratio and 95% confidence interval calculated from Cox proportional hazard models, centering continuous variables and using k-1 indicator variables for ordinal variables with k levels. The assumption of proportional hazards was assessed graphically using log-log plots and Schoenfeld's chi-squared goodness-of-fit procedures²².

We considered the possibility that the method of updating the tumor registry's "date last known alive" with visit data may bias our estimates of survival if one ethnic group was more likely to have contact with the HFMG following diagnosis. Therefore, we conducted the analysis twice: first, we included only tumor registry follow-up dates; second, we used the updated data. We

found negligible differences between the two approaches, therefore, analyses including the updated data are included in this analysis.

Results

We identified 1,321 women members of HAP who were diagnosed with breast cancer between January 1986 through April 1996 and for whom mammography was a fully covered benefit. From this group, we excluded 161 women because they were not assigned to HFMG physicians at the time of diagnosis, and an additional 274 women because they were not continuously enrolled in HAP for one year before diagnosis, for a final sample of 886 women. There was no difference between the study group and the women excluded with respect to the proportion of African Americans (30%).

The median follow-up time was 50 months overall and was similar for African American (49 months for those still alive) and European American women (50 months for those still alive). A total of 137 deaths occurred during the study period. Table 1 shows the baseline demographic and tumor-specific characteristics of the study population. An ordinal logistic regression model indicated that European American women were more likely to have earlier stage disease at diagnosis than African American women ($p=0.007$). Examining this issue more closely, European Americans were more likely than African Americans to have earlier stage (0, I) disease, with a difference of 11% (95% CI 3%, 18%) compared to African Americans. Among those with stage II disease, we found no material difference between African American and European American women in the proportion with lymph node involvement (difference=5%, 95% CI -6%, 17%).

The 5-year survival was 77% for African Americans and 84% for European Americans. The crude estimates, by race are shown in Figure 1. Overall, African Americans had poorer survival compared to European Americans (hazard ratio= 1.6, 95% CI (1.1, 2.2) (Figure 1). There was no evidence of violation of the proportional hazards assumption. Table 2 presents the hazard ratios when we adjusted for stage and sociodemographics, separately and in combination. When stage was added to the model, the hazard ratio decreased to 1.3. Adjusting only for sociodemographics, the hazard ratio was reduced to 1.2. Finally, when we controlled for both stage and sociodemographics, the hazard ratio was reduced to 1.0 (95% CI 0.7, 1.5). The survival curves by race, adjusting for sociodemographic characteristics and stage are shown in Figure 2, and reflect this equivalent survival pattern.

CONCLUSIONS

In a setting with equal access to and delivery of health care, we found that adjustment for sociodemographic variables and stage eliminated the racial differences in breast cancer survival. Sociodemographic variables apparently confounded the association between race and survival more than stage of disease. Our conclusions about the effect of sociodemographic factors is similar to those made by some¹⁹⁻²¹, but not all^{3,6,14,16} studies. Our study adds to the information provided by these others because we included patients within a single HMO and medical group with equal mammography coverage, providing a larger degree of homogeneity in health care access and delivery than in most other studies and perhaps reducing unmeasured differences in confounding factors.

A recent study from the Department of Defense Cancer Tumor Registry found that women with equal access to military care and treatment fared better than the general US population but an unexplained survival difference remained between African American and European American women, even after controlling for stage and other demographic and pathologic factors²³.

However, data on income were not available in this study, however.

The fact that we estimated income from US census data is a limitation of this study. As a result, we expect some degree of misclassification of income. By mapping the addresses to block groups, however, the misclassification should occur to a lesser degree than if we used estimates based on census tracts or zip codes. Our study did not include information on some factors related to survival that also may be related to race, such as treatment and estrogen receptor status.

Nevertheless, we found that in a setting with relatively homogeneous access to health care and treatment, racial differences in survival disappear after adjusting for sociodemographics and stage. Further, census block level income (and what it represents) is a substantial confounder of the race-survival association.

REFERENCES

1. Chevarley F, White E: Recent trends in breast cancer mortality among White and Black US women. *Am J Public Health* 1997;87:775-781.
2. Wingo PA, Ries LAG, Rosenberg HM, Miller DS, Edwards BK: Cancer incidence and mortality, 1973-1995. *Cancer* 1998;82:1197-1207.
3. Bain RP, Greenberg RS, Whitaker JP: Racial differences in survival of women with breast cancer. *J Chron Dis* 1986;39:631-642.
4. Hunter CP, Redmond CK, Chen VW, et al: Breast cancer: factors associated with stage at diagnosis in Black and White women. *J Natl Cancer Inst* 1993;85:1129-1137.
5. West DW, Satariano WA, Ragland DR, Hiatt RA: Comorbidity and breast cancer survival: A comparison between Black and White women. *Ann Epidemiol* 1996;6:413-419.
6. Eley JW, Hill HA, Chen VW, et al: Racial differences in survival from breast cancer. *JAMA* 1994;272:947-954.
7. Burns RB, McCarthy EP, Freund KM, et al: Black women receive less mammography even with similar use of primary care. *Ann Intern Med* 1996;125:173-182.

8. Anderson LM, May DS: Has the use of cervical, breast, and colorectal cancer screening increased in the United States? *Am J Public Health* 1995;85:840-842.
9. Chen VW, Correa P, Kurman RJ, et al: Histological characteristics of breast carcinoma in Blacks and Whites. *Cancer Epidemiol Biomark Prev* 1994;3:127-135.
10. Elledge RM, Clark GM, Chamness GC, Osborne CK: Tumor biologic factors and breast cancer prognosis among White, Hispanic, and Black women in the United States. *J Natl Cancer Inst* 1994;86:705-712.
11. Moormeier J: Breast cancer in Black women. *Ann Intern Med* 1996;124:897-905.
12. Weiss SE, Tartter PI, Ahmed S, et al: Ethnic differences in risk and prognostic factors for breast cancer. *Cancer* 1995;76:268-274.
13. Heimann R, Ferguson D, Powers C, Suri D, Weichselbaum RR, Hellman S: Race and clinical outcome in breast cancer in a series with long-term follow-up evaluation. *J Clin Oncol* 1997;15:2329-2337.
14. Vernon SW, Tilley BC, Neale AV, Steinfeldt L: Ethnicity, survival, and delay in seeking treatment for symptoms of breast cancer. *Cancer* 1985;55:1563-1571.

15. Simons MS, Severson RK: Racial differences in survival of female breast cancer in the Detroit metropolitan area. *Cancer* 1996;77:308-314.
16. Simon MS, Severson RK: Racial differences in breast cancer survival: The interaction of socioeconomic status and tumor biology. *Am J Obstet Gynecol* 1997;176:S223-S239.
17. Perkins P, Cooksley CD, Cox JD: Breast Cancer: is ethnicity an independent prognostic factor for survival? *Cancer* 1996;78:1241-1247.
18. Bassett MT, Krieger N: Social class and Black-White differences in breast cancer survival. *Am J Public Health* 1986;76:1400-1403.
19. Dayal HH, Power RN, Chiu C: Race and socio-economic status in survival from breast cancer. *J Chron Dis* 1982;35:675-683.
20. Gordon NH, Crowe JP, Brumberg DJ, Berger NA: Socioeconomic factors and race in breast cancer recurrence and survival. *Am J Epidemiol* 1992;135:609-618.
21. Ansell D, Whitman S, Lipton R, Cooper R: Race, income, and survival from breast cancer at two public hospitals. *Cancer* 1993;72:2974-2978.
22. Schoenfeld D: Chi-squared goodness-of-fit tests for the proportional hazards regression model. *Biometrika* 1980;67:145-153.

23. Wojcik BE, Spinks MK, Optenberg SA: Breast carcinoma survival analysis for African American and White women in an equal-access health care system. *Cancer* 1998;82:1310-1318.

Table 1. Baseline Demographic and Tumor Characteristics

Variable	African American N = 273	European American N = 613
Sociodemographics		
% Married	(54%)	(59%)
Mean Age (SE) at diagnosis	55 years (\pm 0.8)	56 years (\pm 0.5)
Median household income (SE)	\$26,000 (\pm \$931)	\$44,000 (\pm \$783)
Mean years (SE) HMO enrollment before diagnosis	6.9 years (\pm 0.3)	5.4 years (\pm 0.1)
Tumor Characteristics		
Stage 0	17 %	21 %
I	29 %	36 %
II	40 %	33 %
III	9 %	8 %
IV	5 %	3 %
Mean tumor size (SE)	2.4 (\pm 0.1)	2.1 (\pm 0.1)

Table 2. Effect of Demographic and Tumor Characteristics on Survival Estimates

Variables in Model	Hazard Ratio (African American versus European American)	95% Confidence Interval
Race Only	1.6	(1.1, 2.2)
Race + Stage	1.3	(0.9, 1.9)
Race + Sociodemographics*	1.2	(0.8, 1.9)
Race + Stage + Sociodemographics*	1.0	(0.7, 1.5)

* Age, marital status and median household income

Figure 1. Crude Kaplan-Meier Survival Estimates, by Race





Figure 2. Survival by Race, Adjusted for Age, Income & Marital Status and Stage





DEPARTMENT OF THE ARMY

US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

23 Aug 01

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to the technical reports listed at enclosure. Request the limited distribution statement for these reports be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLIS M. RINEHART
Deputy Chief of Staff for
Information Management

Reports to be Downgraded to Unlimited Distribution

ADB241560	ADB253628	ADB249654	ADB263448
ADB251657	ADB257757	ADB264967	ADB245021
ADB263525	ADB264736	ADB247697	ADB264544
ADB222448	ADB255427	ADB263453	ADB254454
ADB234468	ADB264757	ADB243646	
ADB249596	ADB232924	ADB263428	
ADB263270	ADB232927	ADB240500	
ADB231841	ADB245382	ADB253090	
ADB239007	ADB258158	ADB265236	
ADB263737	ADB264506	ADB264610	
ADB239263	ADB243027	ADB251613	
ADB251995	ADB233334	ADB237451	
ADB233106	ADB242926	ADB249671	
ADB262619	ADB262637	ADB262475	
ADB233111	ADB251649	ADB264579	
ADB240497	ADB264549	ADB244768	
ADB257618	ADB248354	ADB258553	
ADB240496	ADB258768	ADB244278	
ADB233747	ADB247842	ADB257305	
ADB240160	ADB264611	ADB245442	
ADB258646	ADB244931	ADB256780	
ADB264626	ADB263444	ADB264797	